

Recursive sequences

Problem code: HS09REC

Bernie wishes to impress his math teacher with a new theorem. He observes some sequences which satisfy a recursive relation

$$a_{n+2} = 2a_{n+1} - a_n + 2$$

Each sequence of his concern starts with number $a_1 = 1$, but the second numbers differ. Bernie thinks he found a nice rule, which he wants to check. He thinks that no matter what the number a_2 is and no matter which n he chooses, one always can find an element of the sequence which equals $a_n a_{n+1}$.

You can help him in his investigations by finding required elements.

Input

There is K ($1 \leq K \leq 1\,000$) lines of standard input. Each consists of two integer numbers a_2, n ($2 \leq a_2 \leq 1\,000, 1 \leq n \leq 1\,000\,000\,000$) separated by spaces.

The line $K+1$ will contain two zeros, which shouldn't be processed.

Output

Write out K lines of output - one for each testcase. For each testcase the line should contain the smallest positive integer m such that $a_m = a_n a_{n+1}$ or the number 0 if such an m doesn't exist.

Example

Input : 2 1

2 2

2 4

3 5

0 0

Output :

2

4

14

26

Scoring

For solving this problem you will score 10 points.

Added by: Adam Dzedzej

Date: 2010-01-30

Time limit: 1s

Source
limit: 50000B

Languages: SED C99 strict C++ 4.0.0-8 C++ 4.3.2 TCL SCALA NICE NEM BASH PHP SCM guile
LISP sbcl LISP clisp ERL TECS TEXT DOC PDF PS PERL 6 JS

Resource: High School Programming League (thanks to Talent Association)